



Condyloma acuminata on the breast a case report

Dr Birbal kumar, Dr Balkishan Gupta, Dr Deepankar S Mitra

Submitted: 01-01-2023

Accepted: 08-01-2023

ABSTRACT

Rationale: Condyloma acuminatum (CA) is a benign tumor primarily caused by infection with human papillomavirus (HPV) type-6 or type-11, lesions of which are most frequently found on the genital and perianal squamous mucosa and skin. CA outside the genitals is not common.

Patient concerns: A 35-year-old female presented with lesions on the right breast near nipple areola complex and upper inner, lower outer quadrant presenting fungating mass non pedunculated mass with recurrent infection.

Diagnoses: Histopathological examination and HPV detection made a definite diagnosis of CA.

Intervention: The patient was treated with nipple sparing surgical excision of fungating mass followed by circular reduction mammoplasty and topical imiquimod cream.

Outcomes: After 6 months follow-up, there was no sign of recurrence.

Lessons: This case shows that we should pay more attention to CA outside the genitals in the process of diagnosis and treatment.

Keywords: condyloma acuminatum, HPV, nipple, mammoplasty, breast

I. INTRODUCTION

Condyloma acuminatum (CA) is a common sexually transmitted disease among males and females caused by human papillomavirus

(HPV).^[1] HPV is usually cleared within two years of infection, but it is possible for a latency period to occur, with the first occurrence or a recurrence happening months or even years later.^[2] Although, CA can occur on genital and perianal sites through sexual contact, it can also be found outside the genitals by oral-genitals or oral-nipple contact.^[3] Here, we report an uncommon case that a female infected with CA on the right breast

II. CASE REPORT

A 35-year-old female presented with a one year history of some papillary lesions on the right breast. One year ago, the patient noticed a cauliflower lesion on the right breast, which presented reddish, fleshy, no pain, and no itch. Initially, the lesions were ignored because they grew slowly. The patient was otherwise healthy and had history of local infection due to microinjury of fungating mass.

A physical examination 15x3mm cauliflower-like wart was found on the right breast (Fig.1). Surgical excision of fungating mass with circular reduction mammoplasty (Fig.2,3). Histopathological examinations revealed massive epidermal hyperplasia, hyperkeratosis, parakeratosis, and koilocytosis, which were consistent with the diagnosis of CA. Polymerase-chain-reaction assay for HPV DNA revealed an infection with HPV type 11.



Figure 1

Condyloma acuminata on the right breast.



Figure 2
Nipple sparing Surgical excision of fungating mass .



Figure 3
Circular reduction mammoplasty of right breast

Based on the above evidences, a definite diagnosis of CA on the right breast. The patient was treated with surgical removal of fungating mass along with circular reduction mammoplasty and topical imiquimod cream for two weeks. There was no sign of recurrence within one year

III. DISCUSSION

CA is a benign tumor primarily caused by infection with HPV type-6 or type-11.^[4] CA is spread through direct skin-to-skin contact with an infected individual, usually during sex.^[5] Lesions of CA are most frequently found on the genital and perianal squamous mucosa and skin infected



through sexual contact.^[6,7] They can also infect lips, oral cavity, and even other sites outside the genitals through oral-genitals contact. In recent years, some atypical cases of CA occurring at sites like the breast,^[8] tongue,^[9] or groin^[10] have been reported.

We reported the case of CA on the right breast, which shows that in clinical work we should pay more attention to CA outside the genitals, avoiding delay and errors in the process of diagnosis and treatment.

REFERENCES

- [1]. Manini I, Montomoli E. Epidemiology and prevention of human papillomavirus. *Ann Ig*2018;30Suppl 1:28–32.
- [2]. Moscicki AB, Schiffman M, Kjaer S, et al. Chapter 5: Updating the natural history of HPV and anogenital cancer. *Vaccine* 2006;24Suppl 3:S3/42–51.
- [3]. Reis HL, Ferreira DC, Forattini AG, et al. Genital and oral human papillomavirus infection in a patient from the group of women who have sex with women. *Clinics (Sao Paulo, Brazil)*2010;65:1383–5.
- [4]. Chikandiwa A, Kelly H, Sawadogo B, et al. Prevalence, incidence and correlates of low risk HPV infection and anogenital warts in a cohort of women living with HIV in Burkina Faso and South Africa. *PLoS One* 2018;13:e0196018.
- [5]. Doorbar J, Egawa N, Griffin H, et al. Human papillomavirus molecular biology and disease association. *Rev Med Virol* 2015;25Suppl 1:2–3.
- [6]. Yakely AE, Avni-Singer L, Oliveira CR, et al. Human papillomavirus vaccination and anogenital warts: a systematic review of impact and effectiveness in the United States. *Sex Transm Dis*2019;46:213–20.
- [7]. Hamano I, Hatakeyama S, Yamamoto H, et al. Condyloma acuminata of the urethra in a male renal transplant recipient: a case report. *Transplant Proc* 2018;50:2553–7.
- [8]. Saeki Y, Sato S, Okajima K, et al. Condyloma acuminatum of the nipple and areola. *Int J Dermatol* 2014;53:e171–2.
- [9]. Percinoto AC, Danelon M, Crivelini MM, et al. Condyloma acuminata in the tongue and palate of a sexually abused child: a case report. *BMC Res Notes* 2014;7:467.
- [10]. Verma S. Bilateral linear location of condylomata acuminata in the groin: an uncommon presentation aggravated by topical steroids. *Indian J Dermatol* 2014;59:636.