International Journal Dental and Medical Sciences Research
Volume 7, Issue 4, July – Aug. 2025 pp 46-47 www.ijdmsrjournal.com ISSN: 2582-6018

Periodontal Treatment in Epileptic Patients: Clinical Considerations and Management Strategies

Dr. Alejandro Antonio Rivero Garcia

Date of Submission: 20-07-2025 Date of Acceptance: 30-07-2025

ABSTRACT
-Epileptic patients present unique challenges in periodontal care due to the nature of their neurological condition and the side effects of antiepileptic medications. Effective periodontal treatment requires a comprehensive understanding of seizure types, medication-induced gingival overgrowth, and stress-related triggers. This article discusses the clinical management of periodontal disease in epileptic individuals, including preventive strategies, surgical considerations, and safety protocols for in-office care.

I. INTRODUCTION

-Epilepsy is a chronic neurological disorder characterized by recurrent, unprovoked seizures resulting from abnormal electrical activity in the brain. Affecting over 50 million people worldwide, it requires lifelong management with antiepileptic drugs (AEDs). In dental practice, epilepsy is a significant medical consideration, particularly in periodontal therapy, where gingival tissue health and response to treatment are directly influenced by medication and systemic factors.

-The most common oral manifestation related to epilepsy is drug-induced gingival overgrowth, particularly associated with phenytoin, though other AEDs (e.g., valproate, phenobarbital) may also contribute. Moreover, anxiety, stress, and certain dental procedures can potentially trigger seizures during treatment.

Oral Manifestations and Periodontal Risks in Epileptic Patients

-Gingival Overgrowth: Commonly induced by phenytoin; characterized by fibrotic enlargement, primarily in the anterior maxillary and mandibular gingiva.

-Increased Plaque Accumulation: Due to compromised oral hygiene, manual dexterity issues, or cognitive impairments.

-Bleeding and Inflammation: Related to poor plaque control and inflammatory response.

-Dental Trauma: Seizure episodes can result in fractured teeth or soft tissue injuries, complicating periodontal status.

Methods (General Clinical Approach for Periodontal Care)

1. Medical History Review and Seizure Assessment

-Document seizure type, frequency, triggers, and date of last episode

-Identify antiepileptic medications and dosages

-Obtain physician consultation if seizures are poorly controlled

-Determine whether premedication or special precautions are needed

2. Oral Hygiene Instruction and Preventive Care

-Educate the patient and caregivers on plaque control techniques
-Recommend soft toothbrushes or electric brushes, depending on motor skills
-Emphasize the importance of routine maintenance visits
-Use chlorhexidine mouthwash or fluoride rinses if tolerated

3. Non-Surgical Periodontal Therapy

-Scaling and root planing (SRP) to reduce inflammation and control biofilm

-Perform short, stress-free appointments
-Use desensitizing techniques and avoid overstimulation
-Monitor gingival response to improve plaque control before considering surgery

4. Management of Gingival Overgrowth

-If gingival hyperplasia is present:
-Emphasize meticulous oral hygiene to reduce severity
-Collaborate with the physician to consider medication substitution (e.g., switching from phenytoin to carbamazepine if medically appropriate)
-In cases of persistent overgrowth, gingivectomy or periodontal flap surgery may be indicated

5. Surgical Considerations

-Perform surgery only when seizures are well controlled
-Consider the use of local anesthesia with minimal epinephrine to reduce stress and cardiovascular risk
-Schedule treatment early in the day, ideally after meals and medication
-Maintain suction and protective measures to avoid aspiration during a seizure

6. Seizure Precaution During Treatment

-Remove sharp instruments when not in use
-Do not use sedation unless medically cleared
-Keep a padded mouth prop if needed
-Keep the dental chair in a slightly reclined position
-Have emergency protocols and oxygen available

If a seizure occurs:

-Clear the area
-Turn the patient to the side if possible
-Protect the head
-Do not place objects in the mouth during the seizure
-Call emergency services if the seizure lasts more than 5 minutes

II. DISCUSSION

Periodontal treatment in epileptic patients requires thoughtful coordination between dental and medical professionals. Gingival overgrowth not only compromises aesthetics and oral hygiene but can lead to secondary periodontitis if left unmanaged.

Early intervention, preventive strategies, and patient education are essential. In more complex cases, surgical therapy may be necessary, always considering the seizure history and systemic stability of the patient.

III. CONCLUSION

Epileptic patients can receive safe and effective periodontal care when clinicians tailor treatment plans to their medical condition. Preventive maintenance, plaque control, careful monitoring of gingival overgrowth, and well-planned appointments contribute to long-term periodontal health in this medically complex population.

REFERENCES

[1]. Ellis JS, Seymour RA, Steele JG, Robertson P, Butler TJ, Thomason JM. "Prevalence of gingival overgrowth induced by calcium channel blockers: a community-based study." J Clin Periodontol. 1999;26(10):664–667.
[2]. Dongari-Bagtzoglou A. "Drug-associated gingival enlargement." Oral Surg Oral Med Oral Pathol Oral Radiol Endod. 2004;98(4):419–427.
[3]. Ristic L, Vlahovic Svabic M, Sasic S, Zelic O. "The effects of anticonvulsant drugs on the gingiva." Srp Arh Celok Lek. 2005;133(3-4):144–148.