International Journal Dental and Medical Sciences Research
Volume 7, Issue 4, July – Aug. 2025 pp 48-49 www.ijdmsrjournal.com ISSN: 2582-6018

Periodontitis in Diabetic Patients: A Bidirectional Relationship

Dr. Alejandro Antonio Rivero Garcia

Date of Submission: 20-07-2025 Date of Acceptance: 30-07-2025

ABSTRACT
-Periodontitis and diabetes mellitus share a complex, bidirectional relationship. While diabetes increases the risk and severity of periodontal disease, periodontitis may negatively impact glycemic control. Understanding the mechanisms and clinical implications of this association is essential for managing both conditions effectively. This article reviews the pathophysiological link between diabetes and periodontitis, the clinical features, and the principles of periodontal management in diabetic patients.

I. INTRODUCTION

Diabetes mellitus is a chronic systemic condition characterized by hyperglycemia due to insulin deficiency or resistance. Periodontitis is a chronic inflammatory disease affecting the supporting tissues of the teeth, leading to progressive attachment loss and bone destruction.

Multiple studies have confirmed that diabetic patients are more susceptible to periodontal disease, and poorly controlled diabetes is associated with increased periodontal destruction. At the same time, periodontal inflammation may contribute to systemic inflammatory burden, affecting metabolic control.

Diabetes and Periodontal Disease: A Bidirectional Link

How Diabetes Affects Periodontitis

-Impaired Immune Response: Diabetes compromises neutrophil function, impairs chemotaxis and phagocytosis, and enhances susceptibility to infection.
-Altered Collagen Metabolism: Hyperglycemia leads to the formation of advanced glycation end-products (AGEs), which impair collagen repair and increase tissue breakdown.
-Microvascular Changes: Diabetes causes thickening of blood vessel walls, reducing oxygen diffusion and waste removal in periodontal tissues.
-Increased Inflammatory Response: Diabetic individuals show higher levels of pro-inflammatory cytokines (e.g., IL-1β, TNF-α) in gingival tissues and crevicular fluid.

How Periodontitis Affects Diabetes

-Periodontal infection increases systemic inflammation, leading to elevated levels of cytokines that interfere with insulin signaling.
-Periodontal therapy has been shown to improve glycemic control, with studies suggesting a reduction in HbA1c levels by 0.4%–0.6% following treatment.

Clinical Features in Diabetic Patients with Periodontitis

-Increased gingival inflammation, even with similar plaque levels
-Rapid periodontal attachment loss
-More frequent periodontal abscesses
-Delayed healing after periodontal surgery or scaling and root planing
-Higher risk of tooth loss
-These features are more pronounced in poorly controlled diabetic patients (HbA1c >7%).

Methods (General Clinical Approach)

Assessment and Diagnosis

-Full periodontal examination (probing depths, clinical attachment loss, bleeding on probing, mobility)
-Radiographic evaluation for bone loss
-Medical consultation to assess glycemic control (HbA1c, fasting blood glucose)

Classification of diabetes type and duration

Periodontal Therapy in Diabetic Patients

Initial Therapy:

-Oral hygiene instruction
-Scaling and root planing
-Antiseptic rinses (e.g., chlorhexidine)

International Journal Dental and Medical Sciences Research
Volume 7, Issue 4, July – Aug. 2025 pp 48-49 www.ijdmsrjournal.com ISSN: 2582-6018

-Systemic Antibiotics: May be considered in severe cases or for patients with compromised healing
-Surgical Therapy: Periodontal surgery may be indicated after non-surgical therapy; requires stable glycemic control before intervention

-Maintenance Therapy: Regular periodontal maintenance every 3–4 months is crucial for long-term stability

Interdisciplinary Collaboration
-Coordination with the patient's primary care physician or endocrinologist is recommended to monitor and stabilize systemic glucose levels during periodontal treatment.
-Nutritional counseling and lifestyle interventions may also benefit overall outcomes.

Prevention and Patient Education
-Emphasize the importance of oral hygiene in diabetic care plans
-Educate patients about the warning signs of periodontal disease
-Encourage regular dental visits for early detection and intervention
-Support smoking cessation as a key modifier of periodontal risk

II. CONCLUSION

Periodontitis and diabetes are interconnected chronic conditions that influence each other in significant ways. Effective periodontal management in diabetic patients requires a tailored, multidisciplinary approach with a focus on systemic control, professional therapy, and patient education. Treating periodontal inflammation may offer benefits beyond oral health by improving metabolic parameters and enhancing quality of life.

REFERENCES

[1]. Preshaw PM, Alba AL, Herrera D, et al. "Periodontitis and diabetes: a two-way relationship." Diabetologia. 2012;55(1):21–31.
[2]. Taylor GW, Borgnakke WS. "Periodontal disease: associations with diabetes, glycemic control and complications." Oral Dis. 2008;14(3):191–203.